



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**200. Proposed by R. D. CARMICHAEL, Indiana University.**

Find the general solution, in relatively prime integers, of the equation  $x^2 + y^2 = z^4$ .

**201. Proposed by E. T. BELL, Seattle, Washington.**

Eisenstein (*Crelle*, t. 27, p. 282) proposed, as the simplest of several problems: "In the expansion of

$$\frac{1 + z + z^2 + \cdots + z^{p-1}}{(1 - z)^{p-1}} - 1,$$

where  $p$  is prime, to show that the coefficients of the various powers of  $z$  are all divisible by  $p$ ."

**202. Proposed by A. R. SCHWEITZER, Chicago, Ill.**

There exists an infinitude of systems of dyads  $\{\alpha\beta\}$  in 7, 9, 11, etc., elements such that each system has the following properties: (1) if  $\alpha\beta$  is in the set, then  $\beta\alpha$  is not in the set; (2) for each dyad  $\alpha\beta$  in the set there exists an element  $\xi$  such that  $\xi\beta$  and  $\alpha\xi$  are also in the set. For example, such a system is,

12,	23,	34,	45,	56,	67,	78,	89,	91
13,	24,	35,	46,	57,	68,	79,	81,	92
14,	25,	36,	47,	58,	69,	71,	82,	93
51,	62,	73,	84,	95,	16,	27,	38,	49

Investigate the existence of

I. A finite set of triads  $\{\alpha\beta\gamma\}$  such that (1) if  $\alpha\beta\gamma$  is in the set, then  $\beta\gamma\alpha$ ,  $\gamma\alpha\beta$  are also in the set but  $\beta\alpha\gamma$  is not in the set, (2) for each triad  $\alpha\beta\gamma$  in the set there exists an element  $\xi$  such that  $\xi\beta\gamma$ ,  $\alpha\xi\gamma$ ,  $\alpha\beta\xi$  are also in the set.

II. A finite set of tetrads  $\{\alpha\beta\gamma\delta\}$  such that (1) if  $\alpha\beta\gamma\delta$  is in the set, then  $\beta\gamma\alpha\delta$ ,  $\gamma\alpha\beta\delta$ ,  $\gamma\delta\alpha\beta$  are also in the set but  $\beta\alpha\gamma\delta$  is not in the set, (2) for each tetrad  $\alpha\beta\gamma\delta$  in the set there exists an element  $\xi$  such that  $\xi\beta\gamma\delta$ ,  $\alpha\xi\gamma\delta$ ,  $\alpha\beta\xi\delta$ ,  $\alpha\beta\gamma\xi$  are also in the set.

The problem for alternating  $n$ -ads for  $n > 4$  is obvious.

**SOLUTIONS OF PROBLEMS.****ALGEBRA.****382. Proposed by C. E. FLANAGAN, Wheeling, W. Va.**

A few days ago, I deduced the following formula for finding the value of the unknown quantity in a cubic equation having the form,  $x^3 + 3A^2x = B$ .

$$\text{Let } C = \sqrt{\frac{2B}{A^3} + 1} - 1. \text{ Then } x = \frac{2B}{A^2(C^2 + 12)} + \frac{AC}{4}.$$

It is required to show: (1) How this formula was derived; (2) under what conditions does it give exact results; (3) in general, what is its degree of approximation; (4) if possible, modify it so that it will always give exact results; (5) if it cannot be so modified, show why.

**SOLUTION BY THE PROPOSER.**

(1) The formula was derived by equating the volume of a spherical segment of one base and unknown altitude,  $x$ , to  $n$  times the volume of a sphere whose radius is the radius of the base of the spherical segment.

Thus, let  $r$  be the given radius of the base of the spherical segment of one base and  $x$  the unknown altitude. Then we have by hypothesis

$$\frac{1}{3}\pi x^2(3r - x) = n(\frac{4}{3}\pi r^3) = \frac{1}{2}\pi x r^2 + \frac{1}{6}\pi x^3.$$

Hence,

$$3rx^2 - x^3 = 4nr^3,$$

and

$$3r^2x + x^3 = 8nr^3;$$

whence, by adding the two equations and reducing,

$$r^2x + rx^2 = 4nr^3.$$

Whence

$$x = -\frac{r}{2} \pm \frac{r}{2} \sqrt{16n+1}.$$

Letting  $r^2 = A^2$ ,  $B = 8nr^3$ , and  $C = \sqrt{\frac{2B}{A^3} + 1} - 1$ , we have

$$(1) \quad x = \frac{AC}{2}.$$

But

$$\frac{B}{\frac{A^3 C^3}{8} + \frac{3A^3 C}{2}} = 1.$$

Hence,

$$(2) \quad x = \frac{AC}{2} \times \frac{B}{\frac{A^3 C^3}{8} + \frac{3A^3 C}{2}} = \frac{4B}{A^2(C^2 + 12)}.$$

From the last two values of  $x$ , we have, by addition and dividing by 2,

$$(3) \quad x = \frac{2B}{A^2(C^2 + 12)} + \frac{AC}{4}.$$

(2) This always gives exact values when  $n = \frac{1}{2}$ .

As an application, take the cubic equation,

$$x^3 + 12x = 1,120.$$

Here  $A = 2$ ,  $B = 1,120$ , and  $C = \sqrt{281} - 1$ . Hence, from (1),  $x = 15.7$ ; from (2),  $x = 8.4$ , and from (3),  $x = 10.03$ . Thus (3) gives a very close approximation.

*Remark by the Editors.*

The proposer does not answer (3), (4), and (5). The deductions in his solution result from the confusion of letters representing different quantities and using them to represent the same quantity without drawing the necessary conclusion therefrom. Thus, in the expression  $\frac{1}{3}\pi x^2 \times (3r - x)$ , the  $r$  here is the radius of the sphere from which the segment is taken, whereas in  $\frac{1}{2}\pi r^2 x + \frac{1}{6}\pi x^3$ , the  $r$  is the radius of the base of the segment. If these two expressions are equated as they are in the solution above, it follows that  $x = r$  and therefore  $n = \frac{1}{2}$ . It thus follows that when  $n = \frac{1}{2}$  the formula gives correct values for cubics of the proposed form. This seems to us to answer the remaining questions in his problem.

**387. Proposed by H. C. FEEMSTER, York College, Nebraska.**

Sum the following series:

$$(1) \quad \sum_{n=1}^{\infty} \frac{1}{8^{n+1}n!}; \quad (2) \quad \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots 2n+1}{2^{2n+1}(2n-1)!}; \quad (3) \quad \sum_{n=1}^{\infty} \frac{1}{2^{2n}n!}.$$

SOLUTION BY J. A. BULLARD, Worcester Polytechnic Institute.

By Maclaurin's Theorem, we have

$$f(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} D^n f(0), \quad (4)$$

where  $D^n f(0) = [D^n f(x)]_{x=0}$ ,  $0! = 1$  and  $D^0 f(0) = f(0)$ .